

Rapid Urinary Antigen Test in Children with Nasopharyngeal Pneumococcal Carriage

Nazofaringeal Pnömonokok Taşıyıcılığı Olan Sağlıklı Çocuklarda İdrarda Hızlı Antijen Testi Kullanımı

Feyza Koç¹, Zafer Kurugöl¹, Oya Halıcıoğlu², Güldane Koturoğlu¹, Aslı Aslan¹, Şöhret Aydemir³, Sadık Akşit¹

¹Department of Pediatrics, Ege Üniversitesi Faculty of Medicine, İzmir, Turkey

²Pediatrics Clinic, Tepecik Training and Research Hospital, İzmir, Turkey

³Department of Microbiology, Ege University Faculty of Medicine, İzmir, Turkey

Abstract

Objective: To assessment the results of the Binax NOW urinary antigen test in healthy children and the influence of nasopharyngeal pneumococcal carriage on the results of the antigen detection test.

Material and Methods: The study was performed on a total of 223 healthy children aged 2–60 months. All the children enrolled and provided both nasopharyngeal swab specimens for culture and urine samples for the antigen detection test (Binax NOW).

Results: A total of 24 children (11%) were nasopharyngeal carriers of *Streptococcus pneumoniae*. In 16 of these children (66.5%), the pneumococcal urinary antigen test was positive. False positive and false negative rates were found 33.4% and 8.5%, respectively.

Conclusion: It should be considered that a positive Binax NOW test alone is poor for distinguishing the pneumococcal infection from carriage.

(*J Pediatr Inf* 2015; 9: 17-9)

Keywords: Binax NOW, carriage, child, pneumococcal

Özet

Amaç: Bu çalışmanın amacı sağlıklı çocuklarda asemptomatik nazofa, ringeal pnömonokok taşıyıcılığının idrar antijen tarama testi olarak kullanılan Binax NOW testine etkisini araştırmaktır.

Gereç ve Yöntemler: Bu çalışmaya yaşları 2-60 ay arasında değişen 223 sağlıklı çocuk alınmıştır. Bütün çocuklardan boğaz kültür antibiyogramı ve idrar örneği alınmıştır. Alınan idrar örneklerinde pnömonokok antijeni belirlemek üzere Binax NOW (idrar antijen tarama testi) testi yapılmıştır.

Bulgular: Çalışmaya alınan çocukların 24'ünde (%11) nazofaringeal pnömonokok taşıyıcılığı saptanmıştır. Pnömonokok taşıyıcılığı olan çocukların 16'sında (%66,5) idrar pnömonokok antijen tarama testi Binax NOW pozitif saptanmıştır. Binax NOW testinin yalancı pozitiflik oranı %33,4, yalancı negatiflik oranı %8,5 olarak belirlenmiştir.

Sonuç: İdrar pnömonokok antijen tarama testinin pnömonokok hastalıkları dışında sağlıklı çocuklarda pnömonokok taşıyıcılığı olduğunda da pozitif olabileceği akılda tutulmalıdır. (*J Pediatr Inf* 2015; 9: 17-9)

Anahtar kelimeler: Binax NOW, taşıyıcılık, çocuk, pnömonokok

Received/Geliş Tarihi:
13.02.2015

Accepted/Kabul Tarihi:
02.03.2015

Correspondence
Address

Yazışma Adresi:
Feyza Koç, Department of
Pediatrics, Ege
Üniversitesi Faculty of
Medicine,
İzmir, Turkey
Phone: +90 232 390 10 01
E-mail:
feyzaumaykoc@yahoo.com

©Copyright 2015 by Pediatric
Infectious Diseases Society -
Available online at
www.cocukenfeksiyon.org

©Telif Hakkı 2015
Çocuk Enfeksiyon
Hastalıkları Derneği -
Makale metnine
www.cocukenfeksiyon.org
web sayfasından ulaşılabilir.
DOI:10.5152/ced.2015.1970



Introduction

Streptococcus pneumoniae is a major cause of mortality and is the leading cause of bacteremia/sepsis, meningitis, pneumonia, and otitis media in childhood (1). *S. pneumoniae* infections are difficult to accurately diagnose in children. A commercial rapid urinary pneumococcal antigen test (Binax NOW) that detects the C polysaccharide antigen present in all *S. pneumoniae* is used in diagnosis of pneumococcal infection (2). This rapid urinary test has

excellent sensitivity and specificity in adults, but studies performed in children is informed that test's sensitivity and specificity rates is lower than adult's (3, 4). Further, studies report that children with nasopharyngeal carriage of *S. pneumoniae* had high rates of positive test results because of pneumococcal antigen reactions (1, 5). We evaluated the Binax NOW urinary antigen test in healthy children and determined the influence of nasopharyngeal pneumococcal carriage on the results of the antigen detection test.

Material and Methods

The study was performed on a total of 223 healthy children aged 2-60 months and received routine clinical care at Ege University Medicine Faculty Health Child Polyclinic, between September 2009 and March 2010. The study was approved by local ethics committee. This study was conducted in accordance with the principles of the Declaration of Helsinki 2008. Informed consent was obtained from all parents. All children had been vaccinated with 7-valent pneumococcal conjugate vaccine in accordance with national vaccination schedule. All children enrolled and provided both nasopharyngeal swab specimens for culture and urine samples for the antigen detection test. Pneumococci were identified using a visual inspection for typical colony morphology, α -hemolysis, solubility testing, and susceptibility to ethylhydrocupreine (Optochin). Further, the Binax NOW test (*S. pneumoniae* urinary antigen test, Binax, Inc. Scarborough, Maine, USA) was simultaneously conducted for determining the urinary pneumococcal antigen. Binax NOW is a rapid immunochromatographic test and uses a rabbit anti-*S. pneumoniae* antibody that binds to any soluble pneumococcal antigen (C polysaccharide) present in the sample. A swab was dipped into urinary specimen and inserted into the test device. A buffer solution was added, and the device was closed. The results were visually read after 15 min. A pink to purple color on both the sample and control lines indicated a positive result. Statistical analysis was performed with the SPSS for Windows (SPSS for Windows, version 15.0; SPSS Inc., Chicago, IL, USA). The rate of children with and without nasopharyngeal carriage who had Binax NOW test results was compared using the Fisher's exact test. P value <0.05 was considered significant.

Results

Our study included 223 children, ages between 2 and 60 months (23.10 ± 16.98 months, median age=19 months), 134 (60%) males and 89 (40%) females. Pneumococcal nasopharyngeal carriage was present in 24 (10.8%) of 223. In 16 of 24 (%66.7) nasopharyngeal carriage children, pneumococcal antigen was determined positive in their urine samples with the Binax NOW test. Further, in 17 of 199 (8.5%) non-nasopharyngeal pneumococcal carriage children, pneumococcal antigen was determined positive in urine samples (Table 1) ($p < 0.001$). The urine antigen detection test was markedly more likely to indicate a positive result for patients who were nasopharyngeal carriers of pneumococcal than for those who were not.

The sensitivity, specificity, and positive and negative predictive values of the Binax NOW test for the detection of nasopharyngeal carriage of *S. pneumoniae* were 66.6%, 91.5%, 48.5, and 95.8, respectively. False positive and false negative rates were found 33.4% and 8.5% respectively.

Discussion

Our study determined that 24 of 223 (10.8%) children are nasopharyngeal carriers of pneumococci. In some developing countries, rates of pneumococcal carriage in children are 80%-90% (1, 6). This study has been performed in children living in urban areas, and all children were vaccinated with 7-valent pneumococcal conjugate vaccine. We suspect that it was because of these reasons.

We found that rates of positive urinary antigen test results significantly varied according to the status nasopharyngeal carriers of the children (Table 1) ($p < 0.001$). In the study by Hammer et al. (7), they considered the pneumococcal antigen detect with the Binax NOW test in the healthy children who were nasopharyngeal pneumococcal carriers was similarly to our study. As a result of this study, it is emphasized that the test may be useful for the diagnosis of pneumococcal infections with low rates of pneumococcal nasopharyngeal carriage. Other studies mentioned that Binax NOW test did not diagnose pneumococcal infections in children because of a high rate of nasopharyngeal carriage (1, 8). The Binax NOW test was determined positive in 66.5% of children who were nasopharyngeal pneumococcal carriers; however, we found low rates of pneumococcal nasopharyngeal carriage (10.8%) in our study. In contrast, the Binax Now test can lose its specificity in children.

Flores et al. (2) and Moisi et al. (9) mentioned in their study that the Binax Now test can be performed in the children for a rapid diagnosis of *S. pneumoniae* infections. Both studies were performed in the children with invasive pneumococcal infection, and the Binax NOW test was assessed through such specific materials like pleural or cerebrospinal fluid. Further, in these studies, children were not searched if they were nasopharyngeal carriers or not.

Table 1. Streptococcus pneumoniae Binax NOW antigen test and nasopharyngeal culture results.

	Nasopharyngeal carriers	Nasopharyngeal culture (-)	Total
Binax NOW (+)	16	17	33
Binax NOW (-)	8	182	190
Total	24	199	223

$p < 0.001$; Binax NOW (urinary pneumococcal antigen test)

Our study determined that sensitivity, specificity, and positive and negative predictive values of the Binax NOW test for the detection of nasopharyngeal carriage of *S. pneumoniae* were 66.6%, 91.5%, 48.5 and 95.8, respectively. In another study determined that the Binax NOW test from the nasal samples had a sensitivity of 95%, a specificity of 78%, and the positive and negative predictive values were 83% and 93%, respectively (10).

Conclusion

In conclusion, the usefulness of the Binax NOW test can be limited based on the positivity of test in children who are nasopharyngeal carriers. Therefore, we recommend that the clinicians should be careful to consider pneumococcal infection in the presence of rapid pneumococcal urinary antigen positivity alone. In that situation, it can be more sensitive to use other rapid diagnosis tests.

Ethics Committee Approval: Ethics committee approval was received for this study from the local ethics committee.

Informed Consent: Informed consent was obtained from patients and their parents who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - F.K., Z.K.; Design - F.K., Z.K., S.A.; Supervision - O.H., A.A.; Funding - F.K., Z.K., S.A.; Materials - F.K., O.H.; Data Collection and/or Processing - F.K., O.H., Ş.A., G.K.; Analysis and/or Interpretation - F.K., S.A.; Literature Review - F.K., O.H., G.K., A.A.; Writing - F.K., Z.K., O.H., S.A.; Critical Review - Z.K., Ş.A.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Bu çalışma için etik komite onayı yerel etik komite kurulundan alınmıştır.

Hasta Onamı: Yazılı hasta onamı bu çalışmaya katılan hastalardan ve ailelerinden alınmıştır.

Hakem Değerlendirmesi: Dış bağımsız.

Yazar Katkıları: Fikir - F.K., Z.K.; Tasarım - F.K., Z.K.; Denetleme - O.H., A.A.; Veri Toplanması ve/veya İşlemesi - F.K., O.H., Ş.A., G.K.; Analiz ve/veya Yorum - F.K., S.A.; Literatür Taraması - F.K.,

O.H., G.K., A.A.; Yazıyı Yazan - F.K., Z.K., O.H., S.A.; Eleştirel İnceleme - Z.K., Ş.A.

Çıkar Çatışması: Yazarlar çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

References

1. Adegbola RA, Obaro SK, Biney E, Greenwood BM. Evaluation of Binax now Streptococcus pneumoniae urinary antigen test in children in a community with a high carriage rate of pneumococcus. *Pediatr Infect Dis J* 2001; 20: 718-9. [\[CrossRef\]](#)
2. Casado Flores J, Nieto Moro M, Berrón S, Jiménez R, Casal J. Usefulness of pneumococcal antigen detection in pleural effusion for the rapid diagnosis of infection by Streptococcus pneumoniae. *Eur J Pediatr* 2010; 169: 581-4. [\[CrossRef\]](#)
3. Neuman MI, Harper MB. Rapid antigen assay for the diagnosis of pneumococcal bacteremia in children: a preliminary study. *Ann Emerg Med* 2002; 40: 399-404. [\[CrossRef\]](#)
4. Domínguez J, Blanco S, Rodrigo C, et al. Usefulness of urinary antigen detection by an immunochromatographic test for diagnosis of pneumococcal pneumonia in children. *J Clin Microbiol* 2003; 41: 2161-3. [\[CrossRef\]](#)
5. Dowell SF, Garman RL, Liu G, Levine OS, Yang YH. Evaluation of Binax NOW, an assay for the detection of pneumococcal antigen in urine samples, performed among pediatric patients. *Clin Infect Dis* 2001; 32: 824-5. [\[CrossRef\]](#)
6. Mulholland EK, Ogunlesi OO, Adegbola RA, et al. Etiology of serious infections in young Gambian infants. *Pediatr Infect Dis J* 1999; 18: 35-41. [\[CrossRef\]](#)
7. Hamer DH, Egas J, Estrella B, MacLeod WB, Griffiths JK, Sempértegui F. Assessment of the Binax NOW Streptococcus pneumoniae urinary antigen test in children with nasopharyngeal pneumococcal carriage. *Clin Infect Dis* 2002; 34: 1025-8. [\[CrossRef\]](#)
8. Dowell SF, Garman RL, Liu G, Levine OS, Yang YH. Evaluation of Binax NOW, an assay for the detection of pneumococcal antigen in urine samples, performed among pediatric patients. *Clin Infect Dis* 2001; 32: 824-5. [\[CrossRef\]](#)
9. Moisi JC, Saha SK, Falade AG, et al. Enhanced diagnosis of pneumococcal meningitis with use of the Binax NOW immunochromatographic test of Streptococcus pneumoniae antigen: a multisite study. *Clin Infect Dis* 2009; 48: 49-56. [\[CrossRef\]](#)
10. Vuorenoja K, Jalava J, Lindholm L, et al. Detection of Streptococcus pneumoniae carriage by the Binax NOW test with nasal and nasopharyngeal swabs in young children. *Eur J Clin Microbiol Infect Dis* 2012; 31: 703-6. [\[CrossRef\]](#)