

Nail Matrix Arrest during the Course of Hand, Foot, and Mouth Disease: Evaluation of Onychomadesis Cases

Dear Editor,

I read the article by Büyükcım at el. (1) titled "Nail Matrix Arrest during the Course of Hand, Foot, and Mouth Disease: Evaluation of Onychomadesis Cases" with great interest.

Hand, foot, and mouth disease (HFMD) is an infection caused by various enterovirus types affecting especially the children under 5 years of age, mostly children aged between 2-10 (2). Treatment is not required in non-serious cases that are tended to heal spontaneously. HFMD epidemics have increased in recent years and cases with more serious progress have been prevalent (3).

Onychomadesis is characterized by a periodic idiopathic shedding of the nails beginning at the proximal end, possibly caused by the temporary arrest of the function of the nail matrix, and may develop as a result of systemic diseases, malnutrition, trauma, periungual dermatitis, chemotherapy, drugs and infections. Cases emerging 4-8 weeks (4, 5) after the HFMD have been reported. They were found as high as 60% after epidemics (6).

HFMD related cases have been increased in our country since 2011 ,onychomadesis was reported as case reports as was published by Büyükcım et al in your journal (1). Limited number of cases,development of onychomadesis in the late period,admission of the patients to the dermatology policlinics,failure to give detailed information to the patients are some of the reasons of this condition.

Considered from this particular point of view informing the patients with HFMD in regards to onychomadesis, examining the patients 3-4 weeks later and taking detailed history of patients with onychomadesis may determine the true prevalence of the disease.

Ayça Vitrinel, MD

Defne Çöl, MD

E-mail: avitrinel@gmail.com

DOI:10.5152/ced.2015.10



References

1. Büyükcım A, Boyraz MS, Hanalioğlu D, Özsürekcı Y, Cengiz AB, Kara A. Nail Matrix Arrest during the Course of Hand, Foot, and Mouth Disease: Evaluation of Onychomadesis Cases. *J Pediatr Inf* 2015; 9: 64-7 [\[CrossRef\]](#)
2. Haley JC, Hood AF. Hand-foot-and-mouth disease. In: Freedberg IM, Eisen AZ, Wolff K, editors. Fitzpatrick's der-

matology in general medicine. 5th ed. New York: McGraw-Hill; 1999. p. 2403-7.

3. Zhou H, Guo SZ, Zhou H, Zhu YF, Zhang LJ, Zhang W. Clinical characteristics of hand, foot and mouth disease in Harbin and the prediction of severe cases. *Chin Med J (Engl)* 2012; 125: 1261-5.
4. Osterback R, Vuorinen T, Linna M, Susi P, Hyypiä T, Waris M. Coxsackievirus A6 and hand, foot, and mouth disease, Finland. *Emerg Infect Dis* 2009; 15: 1485-8. [\[CrossRef\]](#)
5. Redondo Granado MJ, Torres Hinojal MC, Izquierdo López B. Post viral onychomadesis outbreak in Valladolid. *An Pediatr (Barc)* 2009; 71: 436-9. [\[CrossRef\]](#)
6. Salazar A, Borrás MJ, Cordoba J, et al. Brote de onicomadesis asociado a Síndrome de Boca-Mano-Pie. Onychomadesis outbreak associated with hand, foot and mouth disease. *Boletín Epidemiológico Semanal* 2008; 16: 61-4.
7. Topkarcı Z, Erdoğan B, Yazıcı Z. El-ayak-ağız hastalığının klinik ve demografik özellikleri. *Bakırköy Tıp Dergisi* 2013; 9: 12-5. [\[CrossRef\]](#)

Evaluation of Knowledge, Opinions, and Attitudes of Hospital Staff Except Physicians and Nurses Regarding Seasonal Influenza Vaccine

Dear Editor,

I read the article by Karadağ Öncel et al. (1) "The Evaluation of Knowledge, Opinions and Attitudes of Hospital Staff except Physicians and Nurses about the Seasonal Influenza Vaccine" with great interest (1). Seasonal influenza is a disease that affects people of all ages causing morbidity and mortality especially in immunosuppressive patients and elderly people. Seasonal influenza vaccine is recommended by the ACIP (Advisory Committee on Immunization Practices) and AAP (American Academy of Pediatrics) for all children over 6-month old and all adults (2, 3). Since health care personnel are the medium especially in infecting the disease, they are regarded among the risk groups. In Karadağ Öncel et al.'s (1) study, they found that vaccination rate was very low (18.4%); 54.5% of the vaccinated personnel reported that they had the vaccine not to infect the patients they served for. In the same study, all the study group participants answered the question "I protected the others from influenza when I have the influenza vaccine" 36.9% correctly. This study has demonstrated that the knowledge of hospital staff except physicians and nurses about the seasonal influenza infection and vaccine is below the expected level. In a study done in a pandemic period with the participation of 807 physicians and nurses at Ege University, it was similarly found that the rate of seasonal vaccination rate in the previous year was 19%;